arXiv:0810.5386v1 [math.QA] 29 Oct 2008

Iwahori-Hecke type algebras associated with the Lie superalgebras $A(m,n)$, $B(m,n)$, $C(n)$ and $D(m,n)$

Hiroyuki Yamane

Abstract

In this paper we give Iwahori-Hecke type algebras $H_q(\mathfrak{g})$ associated with the Lie superalgebras $\mathfrak{g} = A(m,n)$, $B(m,n)$, $C(n)$ and $D(m,n)$. We classify the irreducible representations of $H_q(\mathfrak{g})$ for generic q.

Introduction

Recently, motivated by a question posed by V. Serganova [S] and study of the Weyl groupoids [H1][H2] associated with Nichols algebras [AS1][AS2] including generalizations of quantum groups, I. Heckenberger and the author [HY] introduced a notion of 'Coxeter groupoids' (in fact they can be defined as semigroups), and showed that a Matsumoto-type theorem holds for the groupoids, so they have the solvable word problem. We mention that the Coxeter groupoid associated with the affine Lie superalgebra $D^{(1)}(2,1;x)$ was used in the study [HSTY], where Drinfeld second realizations of $U_q(D^{(1)}(2,1;x))$ was analized by physical motivation in recent study of AdS/CFT correspondence.

It would be able to be said that one of the main purposes at present of the representation theory is to study the Kazhdan-Lusztig polynomials (cf. [Hu, 7.9]) and their versions. The polynomials are defined by using the standard and canonical bases of the Iwahori-Hecke algebras. The existence of those bases is closely related to the Matsumoto theorem of the Coxeter groups. So it would be natural to ask what to be the Iwahori-Hecke algebras of the Coxeter groupoids. In this paper, we give a tentative answer to this question for the Coxeter groupoids W associated with the Lie superalgebras $\mathfrak{g} = A(m,n)$, $B(m,n)$, $C(n)$ and $D(m,n)$. We introduce the Iwahori-Hecke type algebra $H_q(\mathfrak{g})$ (in the text, it is also denoted by $H_q(W)$) as q-analogue of the semigroup algebra $\mathbb{C}W/\mathbb{C}0$, where 0 is the zero element of W. We also show that if q is nonzero and not any root of unity, $H_q(\mathfrak{g})$ is semisimple and there exists a natural one-to-one correspondence between the equivalence classes of the irreducible representations of $H_q(\mathfrak{g})$ and those of the Iwahori-Hecke algebra $H_q(W_0)$ associated with the Weyl group W_0 of the Lie algebra $\mathfrak{g}(0)$ obtained as the even part of $\mathfrak{g} = \mathfrak{g}(0) \oplus \mathfrak{g}(1)$.

Until now, no relation has been achieved between the groupoids treated in [SV] and this paper.

This paper is composed of the two sections. Main results and their proofs are given in Section 2. Results of [HY] used in Section 2 are introduced in Section 1.

The author thanks to the referee for careful reading and valuable comments, which encourage him so much to make future study.

1 Preliminary—Matsumoto-type theorem of Coxeter groupoids

This section is preliminary. Here we collect the results which have already been given in [HY] and will be used in the next section.

1.1 Semigroups and Monoids

Let K be a non-empty set. Assume that K has a product map $K \times K \to K$, $(x, y) \mapsto xy$. We call K a *semigroup* if $(xy)z = x(yz)$ for $\forall\, x, y, z \in K$. We call K a *monoid* if K is a semigroup and there exists a unit $1 \in K$, that is, $1x = x1 = x$ for all $x \in K$.

1.2 Free semigroup $F_{-1}(N)$ and Free monoid $F_0(N)$

Let N be a non-empty set. Let $F_{-1}(N)$ be the set of all the finite sequences of elements of N, that is

$$F_{-1}(N) := \coprod_{n=1}^{\infty} N^n = \{(h_1, \dots, h_n) | n \in \mathbb{N}, h_i \in N\}.$$

We regard $F_{-1}(N)$ as the semigroup by

$$(h_1, \dots, h_m)(h_{m+1}, \dots, h_{m+n}) = (h_1, \dots, h_m, h_{m+1}, \dots, h_{m+n}).$$

Then we call $F_{-1}(N)$ a *free semigroup*. Let $F_0(N)$ be the semigroup obtained by adding the unit 1, that is, $F_0(N) := \{1\} \cup F_{-1}(N)$, $1 \notin F_{-1}(N)$, and $1x = x1 = x$ for all $x \in F_0(N)$.

1.3 Semigroup generated by the generators and and defined by the relations

Let $Q = \{(x_j, y_j) | j \in J\}$ be a subset of $F_{-1}(N) \times F_{-1}(N)$, where J is an index set. For g_1, $g_2 \in F_{-1}(N)$, we write $g_1 \sim_1 g_2$ if there exist $j \in J$ and

$(f_1, f_2) \in F_0(N) \times F_0(N)$ such that either of the following (i), (ii), (iii) holds.

(i) $g_1 = f_1 x f_2 \neq g_2 = f_1 y f_2$.
(ii) $g_1 = f_1 y f_2 \neq g_2 = f_1 x f_2$.
(iii) $g_1 = g_2 = f_1 x f_2 = f_1 y f_2$.

For $g,\ g' \in F_{-1}(N)$, we write $g \sim g'$ if $g = g'$ or there exists $r \in \mathbb{N}$ and $g_1, \ldots, g_r \in F_{-1}(N)$ such that $g_1 = g$, $g_r = g'$, and $g_i \sim_1 g_{i+1}$ for $1 \leq i \leq r-1$. Then $F_{-1}(N)/\!\sim$ can be regarded as a semigroup by the product $[g][g'] = [gg']$, where for $g \in F_{-1}(N)$, we denote $[g] := \{g' | g' \sim g\} \in F_{-1}(N)/\!\sim$. We call $F_{-1}(N)/\!\sim$ *the semigroup generated by* N *and defined by the relations* $x_j = y_j$ *(*$j \in J$*).* When there is no fear of misunderstanding, we also denote $[g]$ by its representative g by abuse of notation.

1.4 Free group $F_1(N)$ and Involutive free group $F_2(N)$

Let N be a set. Let N^{-1} be a copy of N so that the bijective map $N \to N^{-1}$, $x \mapsto x^{-1}$, is given. Let $F_1(N)$ be the semigroup generated by

$$\{e\} \cup N \cup N^{-1} \quad \text{(disjoint union)}$$

and defined by the relations

$$ee = ex = xe = e, \quad xx^{-1} = x^{-1}x = e \quad \text{for } \forall x \in N.$$

We call $F_1(N)$ the *free group* over N.

Let $F_2(N)$ be the semigroup generated by

$$\{e\} \cup N \quad \text{(disjoint union)}$$

and defined by the relations

$$ee = ex = xe = e, \quad x^2 = e \quad \text{for } \forall x \in N.$$

We call $F_2(N)$ the *involutive free group* over N. Note that $F_2(N)$ can be identified with the quotient group of $F_1(N)$ in the natural sense:

$$F_2(N) = F_1(N)/\{g_1 x_1^2 g_1^{-1} \cdots g_r x_r^2 g_r^{-1} | r \in \mathbb{N}, x_i \in N, g_i \in F_1(N)\}.$$

1.5 Action $\triangleright$ of $F_2(N)$ on A

Let N and A be non-empty sets. An action $\triangleright$ of $F_2(N)$ on A is a map

$$\triangleright : F_2(N) \times A \to A$$

such that

$$e \triangleright a = a, \quad g \triangleright (h \triangleright a) = (gh) \triangleright a \quad \text{for } \forall g, \forall h \in F_2(N), \forall a \in A.$$

Note that $n \triangleright (n \triangleright a) = a$ for all $n \in N, a \in A$.

For n, $n' \in N$ and $a \in A$, define

$$\Theta(n, n'; a) := \{(nn')^m \triangleright a,\ (n'n)^m \triangleright a \,|\, m \in \mathbb{N} \cup \{0\}\}.$$

Let

$$\theta(n, n'; a) := |\Theta(n, n'; a)|.$$

This is the cardinality of $\Theta(n, n'; a)$, which is either in $\mathbb{N}$ or is ∞. One obviously has $\Theta(n, n'; a) = \Theta(n', n; a)$ and $\Theta(n, n'; n \triangleright a) = n \triangleright \Theta(n', n; a)$.

Let $a_0 := a$, $b_0 := a$, and define recursively $a_{m+1} := n \triangleright b_m$, $b_{m+1} := n' \triangleright a_m$ for all $m \in \mathbb{N} \cup \{0\}$. That is:

$$\begin{array}{llll} b_0 := a, & a_1 := n \triangleright a, & b_2 := n' \triangleright n \triangleright a, & a_3 := n \triangleright n' \triangleright n \triangleright a, \ldots \\ a_0 := a, & b_1 := n' \triangleright a, & a_2 := n \triangleright n' \triangleright a, & b_3 := n' \triangleright n \triangleright n' \triangleright a, \ldots \end{array}$$

Then we have

$$\theta(n, n'; a) = \begin{cases} \infty & \text{if } a_m \neq b_m \text{ for all } m \in \mathbb{N}, \\ \min\{m \in \mathbb{N} \,|\, a_m = b_m\} & \text{otherwise.} \end{cases}$$

1.6 Coxeter groupoids

Definition 1.1. [HY, Definition 1] Let N and A be non-empty sets. Let $\triangleright$ be a transitive action of $F_2(N)$ on A. For each $a \in A$ and $i, j \in N$ with $i \neq j$ let

$$m_{i,j;a} = m_{j,i;a} \in (\mathbb{N} + 1) \cup \{\infty\}$$

be such that $\theta(i, j; a) \in \mathbb{N} \Longrightarrow \frac{m_{i,j;a}}{\theta(i,j;a)} \in \mathbb{N} \cup \{\infty\}$ or $\theta(i, j; a) = \infty \Longrightarrow m_{i,j;a} = \infty$. Set

$$\mathbf{m} := (m_{i,j;a} \,|\, i, j \in N, i \neq j, a \in A).$$

Let

$$W \quad = \quad (W, N, A, \triangleright, \mathbf{m}) \tag{1}$$

be the semigroup generated by the set

$$\{0, e_a, s_{i,a} \,|\, a \in A, i \in N\}$$

and defined by the relations

$$00 = e_a 0 = 0 e_a = s_{i,a} 0 = 0 s_{i,a} = 0, \tag{2}$$

$$\begin{aligned} e_a^2 &= e_a, & e_a e_b &= 0 \text{ for } a \neq b, \\ e_{i \triangleright a} s_{i,a} &= s_{i,a} e_a = s_{i,a}, & s_{i, i \triangleright a} s_{i,a} &= e_a, \end{aligned} \tag{3}$$

$$\begin{aligned} &s_i s_j \cdots s_j s_{i,a} = s_j s_i \cdots s_i s_{j,a} \ (m_{i,j;a} \text{ factors}) && \text{if } m_{i,j;a} \text{ is finite and odd,} \\ &s_j s_i \cdots s_j s_{i,a} = s_i s_j \cdots s_i s_{j,a} \ (m_{i,j;a} \text{ factors}) && \text{if } m_{i,j;a} \text{ is finite and even,} \end{aligned} \tag{4}$$

where we use the convention:

$$s_j s_{i,a} := s_{j, i \triangleright a} s_{i,a}, \quad s_i s_j s_{i,a} := s_{i, j \triangleright i \triangleright a} s_j s_{i,a}, \ldots \tag{5}$$

See also (7) below.

1.7 Sign representation

Let $\mathbb{Z}A$ be the free $\mathbb{Z}$-module generated by A, that is,

$$\mathbb{Z}A = \oplus_{a \in A} \mathbb{Z}a.$$

Then there exists a unique semigroup homomorphism

$$\widetilde{\operatorname{sgn}} : W \to \operatorname{End}_{\mathbb{Z}}(\mathbb{Z}A)$$

such that

$$\widetilde{\operatorname{sgn}}(0)(b) = 0, \qquad \widetilde{\operatorname{sgn}}(e_a)(b) = \delta_{ab} b, \qquad \widetilde{\operatorname{sgn}}(s_{i,a})(b) = (-1) \delta_{ab} i \triangleright a \tag{6}$$

for $a, b \in A$ and $i \in N$, where δ means Kronecker's symbol. Hence for $w \in W$ one has

$$w \neq 0$$

if and only if $w = e_a$ for some $a \in A$ or there exist $m \in \mathbb{N}$ and $i_j \in N$, $b_j \in A$ with $1 \leq j \leq m$ such that $b_j = i_{j+1} \triangleright b_{j+1}$ and $w = s_{i_1, b_1} \cdots s_{i_{m-1}, b_{m-1}} s_{i_m, b_m}$. If this is the case, we use the convention

$$s_{i_1} \cdots s_{i_{m-1}} s_{i_m, b_m} := w, \tag{7}$$

and, if $m = 0$, $s_{i_1} \cdots s_{i_{m-1}} s_{i_m, a}$ means e_a. We note again

Lemma 1.2. (1) $s_{i_1} \cdots s_{i_{m-1}} s_{i_m, a} \neq 0$ *for all* $a \in A$ *and* $m \in \mathbb{N} \cup \{0\}$.

(2) *If* $s_{i_1} \cdots s_{i_{m-1}} s_{i_m, a} = s_{j_1} \cdots s_{j_{r-1}} s_{j_r, b}$, *then* $a = b$, $i_1 \cdots i_m \triangleright a = j_1 \cdots j_r \triangleright b$ *and* $(-1)^m = (-1)^r$.

1.8 Generalization of Root systems

Definition 1.3. [HY, Definition 2] We call a quadruple $(R, N, A, \triangleright)$ a *multi-domains root system* if the following conditions hold.

1. N and A are non-empty sets and $\triangleright$ is a transitive action of $F_2(N)$ on A.
2. Let V_0 be the $|N|$-dimensional $\mathbb{R}$-linear space. Then
$$R = \{(R_a, \pi_a, S_a) \,|a \in A\},$$
where $\pi_a = \{\alpha_{n,a} \,|\, n \in N\} \subset R_a \subset V_0$, and π_a is a basis of V_0 for all $a \in A$.
3. $R_a = R_a^+ \cup -R_a^+$ for all $a \in A$, where $R_a^+ = (\mathbb{N} \cup \{0\})\pi_a \cap R_a$.
4. For any $i \in N$ and $a \in A$ one has $\mathbb{R}\alpha_{i,a} \cap R_a = \{\alpha_{i,a}, -\alpha_{i,a}\}$.
5. $S_a = \{\sigma_{i,a} \,|\, i \in N\}$, and for each $a \in A$ and $i \in N$ one has $\sigma_{i,a} \in \mathrm{GL}(V_0)$,
$$\sigma_{i,a}(R_a) = R_{i \triangleright a}, \quad \sigma_{i,a}(\alpha_{i,a}) = -\,\alpha_{i,i\triangleright a}, \quad \sigma_{i,a}(\alpha_{j,a}) \in \alpha_{j,i\triangleright a} + (\mathbb{N} \cup \{0\})\alpha_{i,i\triangleright a}$$
for all $j \in N \setminus \{i\}$.
6. $\sigma_{i,i\triangleright a}\sigma_{i,a} = \mathrm{id}$ for $a \in A$ and $i \in N$.
7. Let $a \in A$, $i, j \in N$, $i \neq j$, and $d = |((\mathbb{N} \cup \{0\})\alpha_{i,a} + (\mathbb{N} \cup \{0\})\alpha_{j,a}) \cap R_a|$. If d is finite then $\theta(i, j; a)$ is finite and it divides d.

Convention. We write
$$(R, N, A, \triangleright) \in \mathcal{R}$$
if $(R, N, A, \triangleright)$ is a multi-domains root system, that is, $\mathcal{R} = \{(R, N, A, \triangleright)\}$ denotes the family of all the multi-domains root systems.

Definition 1.4. [HY, Definition 4] Let $(R, N, A, \triangleright) \in \mathcal{R}$. Let $\mathbf{m} := (m_{i,j;a} \,|\, i, j \in N, i \neq j, a \in A)$ be such that $m_{i,j;a} := |((\mathbb{N} \cup \{0\})\alpha_{i,a} + (\mathbb{N} \cup \{0\})\alpha_{j,a}) \cap R_a|$. Then we call $(W, N, A, \triangleright, \mathbf{m})$ the *Coxeter groupoid associated with* $(R, N, A, \triangleright)$.

Theorem 1.5. [HY, Theorem 1] *Let $(R, N, A, \triangleright) \in \mathcal{R}$. Set $V = \bigoplus_{a \in A} V_a$, where $V_a = V_0$. Let $P_a : V \to V_a$ and $\iota_a : V_a \to V$ be the canonical projection and the canonical inclusion map respectively. Then the assignment $\rho : 0 \mapsto 0 \cdot \mathrm{id}_V$, $e_a \mapsto \iota_a P_a$, $s_{i,a} \mapsto \iota_{i\triangleright a}\sigma_{i,a}P_a$, gives a faithful representation (ρ, V) of the Coxeter groupoid $(W, N, A, \triangleright, \mathbf{m})$ associated with $(R, N, A, \triangleright)$.*

1.9 Matsumoto-type theorem

Define $\ell : W \to \mathbb{N} \cup \{0\} \cup \{-\infty\}$ to be the map such that $\ell(0) = -\infty$, $\ell(e_a) = 0$ for all $a \in A$, and

$$\ell(w) = \min\{m \in \mathbb{N} \,|\, w = s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a} \text{ for some } i_1, \ldots, i_m \in N,\, a \in A\}$$

for all $w \in W \setminus (\{0\} \cup \{e_a | a \in A\})$; we also refer to Lemma 1.2 (1) for this definition of ℓ. One has

$$\ell(w) = \ell(w^{-1}) \tag{8}$$

for $w \in W \setminus \{0\}$, and

$$\ell(ww') \leq \ell(w) + \ell(w') \tag{9}$$

for $w, w' \in W$. We say that a product $w = s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a} \in W$ is *reduced* if $m = \ell(w)$.

Definition 1.6. [HY, Definition 5] Let $W = (W, N, A, \triangleright, \mathbf{m})$ be a Coxeter groupoid. Let $\widetilde{W} = (\widetilde{W}, N, A, \triangleright, \mathbf{m})$ denote the semigroup generated by the set $\{0, \tilde{e}_a, \tilde{s}_{i,a} \,|\, a \in A, i \in N\}$ and defined by the relations

$$00 = 0, \quad 0\tilde{e}_a = \tilde{e}_a 0 = 0\tilde{s}_{i,a} = \tilde{s}_{i,a}0 = 0, \tag{10}$$

$$\tilde{e}_a^2 = \tilde{e}_a, \quad \tilde{e}_a\tilde{e}_b = 0 \text{ for } a \neq b, \quad \tilde{e}_{i\triangleright a}\tilde{s}_{i,a} = \tilde{s}_{i,a}\tilde{e}_a = \tilde{s}_{i,a}, \tag{11}$$

$$\begin{array}{ll} \tilde{s}_i\tilde{s}_j \cdots \tilde{s}_j\tilde{s}_{i,a} = \tilde{s}_j\tilde{s}_i \cdots \tilde{s}_i\tilde{s}_{j,a} \ (m_{i,j;a} \text{ factors}) & \text{if } m_{i,j;a} \text{ is finite and odd,} \\ \tilde{s}_j\tilde{s}_i \cdots \tilde{s}_j\tilde{s}_{i,a} = \tilde{s}_i\tilde{s}_j \cdots \tilde{s}_i\tilde{s}_{j,a} \ (m_{i,j;a} \text{ factors}) & \text{if } m_{i,j;a} \text{ is finite and even.} \end{array} \tag{12}$$

Theorem 1.7. [HY, Theorem 5] (Matsumoto-type theorem of the Coxeter groupoids) *Let $W = (W, N, A, \triangleright, \mathbf{m})$ be the Coxeter groupoid associated with $(R, N, A, \triangleright) \in \mathcal{R}$ (see Definition 1.4). Suppose that $m \in \mathbb{N} \cup \{0\}$, $a \in A$, and $(i_1, \ldots, i_m), (j_1, \ldots, j_m) \in N^m$ such that $\ell(s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a}) = m$ and equation*

$$s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a} = s_{j_1} \cdots s_{j_{m-1}} s_{j_m,a}$$

holds in W. Then in the semigroup $(\widetilde{W}, N, A, \triangleright, \mathbf{m})$ one has

$$\tilde{s}_{i_1} \cdots \tilde{s}_{i_{m-1}} \tilde{s}_{i_m,a} = \tilde{s}_{j_1} \cdots \tilde{s}_{j_{m-1}} \tilde{s}_{j_m,a}.$$

Corollary 1.8. [HY, Corollary 6] *Let $W = (W, N, A, \triangleright, \mathbf{m})$ be the Coxeter groupoid associated with $(R, N, A, \triangleright) \in \mathcal{R}$ (see Definition 1.4). Suppose that $m \in \mathbb{N} \cup \{0\}$, $a \in A$, and $(i_1, \ldots, i_m) \in N^m$ such that $\ell(s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a}) < m$ holds in $(W, N, A, \triangleright, \mathbf{m})$. Then there exist $j_1, \ldots, j_m \in N$ and $t \in \{1, \ldots, m-1\}$ such that $j_t = j_{t+1}$ and in the semigroup $(\widetilde{W}, N, A, \triangleright, \mathbf{m})$ one has the equation*

$$\tilde{s}_{i_1} \cdots \tilde{s}_{i_{m-1}} \tilde{s}_{i_m,a} = \tilde{s}_{j_1} \cdots \tilde{s}_{j_t} \tilde{s}_{j_{t+1}} \cdots \tilde{s}_{j_{m-1}} \tilde{s}_{j_m,a}.$$



Figure 1: Dynkin diagrams of the Lie superalgebra $D(2,1;x)$

In the next section, we also need

Proposition 1.9. [HY, Corollary 3] *Let* $m \in \mathbb{N}$*,* $(i_1,\ldots,i_m,j) \in N^{m+1}$*, and* $a \in A$*, and suppose that* $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}) = m$*. Then:*

(1) $m = |\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(R^+_a) \cap -R^+_{i_1\cdots i_m\triangleright a}|$.

(2) $\ell(s_{i_1}\cdots s_{i_m}s_{j,j\triangleright a}) = m-1 \iff \sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j,a}) \in -R^+_{i_1\cdots i_m\triangleright a}$.

(3) $\ell(s_{i_1}\cdots s_{i_m}s_{j,j\triangleright a}) = m+1 \iff \sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j,a}) \in R^+_{i_1\cdots i_m\triangleright a}$.

Example 1.10. Here we treat the finite dimensional simple Lie superalgebra $D(2,1;x)$, where $x \notin \{0,-1\}$. Note that it has 14 (positive and negative) roots. One has $m_{i,j;a(k)} = 2 + \delta_{k0} + (1-\delta_{k0})(\delta_{ik}+\delta_{jk})$ and $\sigma_{i,a(k)}(\alpha_{j,a(k)}) = \alpha_{j,i\triangleright a(k)} + \delta_{3,m_{i,j;a(k)}}\alpha_{i,i\triangleright a(k)}$ for $i \neq j$. Moreover

$$\begin{aligned} R^+_{a(k)} = \ & \pi_{a(k)} \cup \{\alpha_{i,a(k)} + \alpha_{j,a(k)} | m_{i,j;a(k)} = 3\} \\ & \cup \{\alpha_{1,a(k)} + \alpha_{2,a(k)} + \alpha_{3,a(k)}\} \\ & \cup \{\alpha_{i,a(k)} + 2\alpha_{k,a(k)} + \alpha_{j,a(k)} | m_{i,j;a(k)} = 2\}. \end{aligned}$$

Note that $D(2,1;1) = D(2,1) = \mathfrak{osp}(4|2)$ (see also Section 2.2). Let $w_{a(2)} := s_{3,a(2)}s_{2,a(0)}s_{3,a(3)}s_{1,a(3)}s_{3,a(0)}s_{2,a(2)}s_{1,a(2)}$. Then $\rho(w_{a(2)}) = -\mathrm{id}_{V_{a(2)}}$. Indeed:

$$\begin{aligned} &\alpha_{1,a(2)} \mapsto -\alpha_{1,a(2)} \mapsto -\alpha_{1,a(0)} - \alpha_{2,a(0)} \mapsto -\alpha_{1,a(3)} - \alpha_{2,a(3)} - 2\alpha_{3,a(3)} \\ &\quad \mapsto -\alpha_{1,a(3)} - \alpha_{2,a(3)} - 2\alpha_{3,a(3)} \mapsto -\alpha_{1,a(0)} - \alpha_{2,a(0)} \mapsto -\alpha_{1,a(2)} \mapsto -\alpha_{1,a(2)}, \\ &\alpha_{2,a(2)} \mapsto \alpha_{1,a(2)} + \alpha_{2,a(2)} \mapsto \alpha_{1,a(0)} \mapsto \alpha_{1,a(3)} + \alpha_{3,a(3)} \mapsto \alpha_{3,a(3)} \mapsto -\alpha_{3,a(0)} \\ &\quad \mapsto -\alpha_{2,a(2)} - \alpha_{3,a(2)} \mapsto -\alpha_{2,a(2)}, \\ &\alpha_{3,a(2)} \mapsto \alpha_{3,a(2)} \mapsto \alpha_{2,a(0)} + \alpha_{3,a(0)} \mapsto \alpha_{2,a(3)} \mapsto \alpha_{2,a(3)} \mapsto \alpha_{2,a(0)} + \alpha_{3,a(0)} \\ &\quad \mapsto \alpha_{3,a(2)} \mapsto -\alpha_{3,a(2)}. \end{aligned}$$

By Proposition 1.9(1), we have $\ell(w_{a(2)}) = |R^+_{a(2)}| = 7$, $w_{a(2)}$ is the longest word. Let $w' := (s_{3,a(2)}s_{2,a(0)})^{-1}w_{a(2)}$. Then $\ell(w') = 5$. By Theorem 1.7, w' has the following four reduced expressions:

$$\begin{aligned} w' =& \underline{s_{3,a(3)}s_{1,a(3)}s_{3,a(0)}}s_{2,a(2)}s_{1,a(2)} = s_{1,a(1)}s_{3,a(1)}\underline{s_{1,a(0)}s_{2,a(2)}s_{1,a(2)}} \\ =& s_{1,a(1)}\underline{s_{3,a(1)}s_{2,a(1)}}s_{1,a(0)}s_{2,a(2)} = s_{1,a(1)}s_{2,a(1)}s_{3,a(1)}s_{1,a(0)}s_{2,a(2)}. \end{aligned}$$

2 Main theorems—Irreducible representations of the Iwahori-Hecke type algeras $H_q(A(m,n))$, $H_q(B(m,n))$, $H_q(C(n))$ and $H_q(D(m,n))$ associated with the Lie superalgebras $A(m,n)$, $B(m,n)$, $C(n)$, $D(m,n)$

2.1 Definition of Lie superalgebras

As for the terminology concerning Lie superalgebras, we refer to [K].

Let $\mathfrak{v} = \mathfrak{v}(0) \oplus \mathfrak{v}(1)$ be a $\mathbb{Z}/2\mathbb{Z}$-graded $\mathbb{C}$-linear space. If $i \in \{0,1\}$ and $j \in \mathbb{Z}$ such that $j - i \in 2\mathbb{Z}$ then let $\mathfrak{v}(j) = \mathfrak{v}(i)$. If $X \in \mathfrak{v}(0)$ (resp. $X \in \mathfrak{v}(1)$) then we write

$$\deg(X) = 0 \text{ (resp. } \deg(X) = 1) \tag{13}$$

and we say that X is an *even* (resp. *odd*) element. If $X \in \mathfrak{v}(0) \cup \mathfrak{v}(1)$, then we say that X is a *homogeneous* element and that $\deg(X)$ is the *parity* (or *degree*) of X. If $\mathfrak{w} \subset \mathfrak{v}$ is a subspace and $\mathfrak{w} = (\mathfrak{w} \cap \mathfrak{v}(0)) \oplus (\mathfrak{w} \cap \mathfrak{v}(1))$ (resp. $\mathfrak{w} \subset \mathfrak{v}(0)$, resp. $\mathfrak{w} \subset \mathfrak{v}(1)$), then we say that $\mathfrak{w}$ is a *graded* (resp. *even*, resp. *odd*) subspace.

Let $\mathfrak{g} = \mathfrak{g}(0) \oplus \mathfrak{g}(1)$ be a $\mathbb{Z}/2\mathbb{Z}$-graded $\mathbb{C}$-linear space equipped with a bilinear map $[\,,\,] : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{g}$ such that $[\mathfrak{g}(i), \mathfrak{g}(j)] \subset \mathfrak{g}(i+j)$ (i, $j \in \mathbb{Z}$); we recall from the above paragraph that

$$\mathfrak{g}(i) = \{X \in \mathfrak{g} \,|\, \deg(X) = i\}. \tag{14}$$

We say that $\mathfrak{g} = (\mathfrak{g}, [\,,\,])$ is a ($\mathbb{C}$-)*Lie superalgebra* if for all homogeneous elements X, Y, Z of $\mathfrak{g}$ the following equations hold.

$$\begin{aligned} [Y,X] =& -(-1)^{\deg(X)\deg(Y)}[X,Y], && \text{(skew-symmetry)} \\ [X,[Y,Z]] =& [[X,Y],Z] + (-1)^{\deg(X)\deg(Y)}[Y,[X,Z]]. && \text{(Jacobi identity)} \end{aligned}$$

We call the Lie algebra $\mathfrak{g}(0)$ the *even part* of $\mathfrak{g}$.

2.2 Lie superalgebras $\mathfrak{gl}(m+1|n+1)$ and $\mathfrak{osp}(m|n)$

Let m, $n \in \mathbb{N} \cup \{0\}$. Let:

$$\mathcal{D}_{m+1|n+1} := \{(p_1, \ldots, p_{m+n+2}) \in \mathbb{Z}^{m+n+2} \,|\, p_i \in \{0,1\}, \sum_{i=1}^{m+n+2} p_i = n+1\}.$$

For $i, j \in \{1, \ldots, m+n+2\}$, let $\mathbf{E}_{i,j}$ denote the $(m+n+2) \times (m+n+2)$ matrix having 1 in (i,j) position and 0 otherwise, that is, the (i,j)-matrix unit. Let $\mathbf{E}_{m+n+2}$ denote the $(m+n+2) \times (m+n+2)$ unit matrix, that is, $\mathbf{E}_{m+n+2} = \sum_{i=1}^{m+n+2} \mathbf{E}_{i,i}$. Denote by $\mathrm{M}_{m+n+2}(\mathbb{C})$ the $\mathbb{C}$-linear space of the $(m+n+2) \times (m+n+2)$-matrices, i.e., $\mathrm{M}_{m+n+2}(\mathbb{C}) = \oplus_{i,j=1}^{m+n+2} \mathbb{C}\mathbf{E}_{i,j}$.

Let $d = (p_1, \ldots, p_{m+n+2}) \in \mathcal{D}_{m+1|n+1}$. The Lie superalgebra $\mathfrak{gl}(m+1|n+1) = \mathfrak{gl}(d)$ is defined by $\mathfrak{gl}(d) = \mathrm{M}_{m+n+2}(\mathbb{C})$ (as a $\mathbb{C}$-linear space),

$$\mathfrak{gl}(d)(0) = \oplus_{1 \leq p_i = p_j \leq m+n+2} \mathbb{C}\mathbf{E}_{i,j}, \quad \mathfrak{gl}(d)(1) = \oplus_{1 \leq p_i \neq p_j \leq m+n+2} \mathbb{C}\mathbf{E}_{i,j}, \tag{15}$$

and $[X, Y] = XY - (-1)^{r_1 r_2} YX$ for $X \in \mathfrak{gl}(d)(r_1)$ and $Y \in \mathfrak{gl}(d)(r_2)$,

where XY and YX mean the matrix product, that is, $\mathbf{E}_{i,j}\mathbf{E}_{k,l} = \delta_{j,k}\mathbf{E}_{i,l}$. Define the $\mathbb{C}$-linear map $\mathrm{str} : \mathfrak{gl}(d) \to \mathbb{C}$ by $\mathrm{str}(\mathbf{E}_{i,j}) = \delta_{i,j}(-1)^{p_i}$. The Lie subsuperalgebra $\{X \in \mathfrak{gl}(d) \,|\, \mathrm{str}(X) = 0\}$ of $\mathfrak{gl}(d)$ is denoted as $\mathfrak{sl}(m+1|n+1) = \mathfrak{sl}(d)$. The finite dimensional simple Lie superalgebra $A(m,n)$ is defined as follows. Let $\mathfrak{z}$ be the one dimensional ideal $\mathbb{C}\mathbf{E}_{m+n+2}$ of $\mathfrak{gl}(d)$. If $m \neq n$, then $A(m,n)$ means $\mathfrak{sl}(d)$. On the other hand, $A(n,n)$ means $\mathfrak{sl}(d)/\mathfrak{z}$, and is also denoted as $\mathfrak{psl}(n+1|n+1)$.

Let $d = (p_1, \ldots, p_{m+2n}) \in \mathcal{D}_{m|2n}$. Define the map $\theta : \{1, \ldots, m+2n\} \to \{1, \ldots, m+2n\}$ by $\theta(i) = m+2n+1-i$. Assume that $p_{\theta(i)} = p_i$. Let $g_i \in \{1, -1\}$ be such that $g_i = -1$ if $p_i = 1$ and $i < \theta(i)$ and $g_i = 1$ otherwise. We have an automorphism Ω of $\mathfrak{gl}(d)$ defined by $\Omega(\mathbf{E}_{i,j}) = -(-1)^{p_i p_j + p_j} g_i g_j \mathbf{E}_{\theta(j),\theta(i)}$. The Lie superalgebra $\mathfrak{osp}(m|2n)$ means $\{X \in \mathfrak{gl}(d) | \Omega(X) = X\}$. We also denote $\mathfrak{osp}(m|2n)$ as follows:

$$\begin{aligned} B(m-1,n) &= \mathfrak{osp}(2m-1|2n) && \text{if } m,\, n \in \mathbb{N}, \\ D(m+1,n) &= \mathfrak{osp}(2m+2|2n) && \text{if } m,\, n \in \mathbb{N}, \\ C(n+1) &= \mathfrak{osp}(2|2n) && \text{if } n \in \mathbb{N}. \end{aligned}$$

We also note that $\mathfrak{osp}(2m+1|0)$, $\mathfrak{osp}(0|2n)$, and $\mathfrak{osp}(2m|0)$ are isomorphic to the simple Lie algebras of type B_m (if $m \geq 2$), C_n (if $n \geq 3$) and D_m (if $m \geq 4$) respectively, so $\mathfrak{osp}(2m+1|0) \cong \mathfrak{o}_{2m+1}$, $\mathfrak{osp}(0|2n) \cong \mathfrak{sp}_{2n}$ and $\mathfrak{osp}(2m|0) \cong \mathfrak{o}_{2m}$. As for the even part $\mathfrak{osp}(m|2n)(0)$ of $\mathfrak{osp}(m|2n)$, we have

$$\mathfrak{osp}(m|2n)(0) \cong \mathfrak{osp}(m|0) \oplus \mathfrak{osp}(0|2n). \tag{16}$$

2.3 Definition of Iwahori-Hecke type algebras

Definition 2.1. Let $W = (W, N, A, \triangleright, \mathbf{m})$ be the groupoid introduced in (1). Assume that A is finite. Let $q \in \mathbb{C}$. Let $H_q(W)$ be the $\mathbb{C}$-algebra (with 1) generated by

$$\{E_a, T_{i,a} | a \in A, i \in N\} \tag{17}$$

and defined by the relations

$$E_a^2 = E_a, \tag{18}$$

$$E_{i \triangleright a} T_{i,a} E_a = T_{i,a}, \tag{19}$$

$$\sum_{a \in A} E_a = 1 \tag{20}$$

$$E_a E_b = 0 \qquad \text{if } a \neq b, \tag{21}$$

$$(T_{i,a} - qE_a)(T_{i,a} + E_a) = 0 \qquad \text{if } i \triangleright a = a, \tag{22}$$

$$T_{i, i \triangleright a} T_{i,a} = E_a \qquad \text{if } i \triangleright a \neq a, \tag{23}$$

$$T_i T_j \cdots T_j T_{i,a} = T_j T_i \cdots T_i T_{j,a} \ (m_{i,j;a} \text{ factors}) \quad \text{if } m_{i,j;a} \text{ is finite and odd,} \tag{24}$$

$$T_j T_i \cdots T_j T_{i,a} = T_i T_j \cdots T_i T_{j,a} \ (m_{i,j;a} \text{ factors}) \quad \text{if } m_{i,j;a} \text{ is finite and even,} \tag{25}$$

where, in (24)-(25), we use the same convention as that of (5) with $s_{i,a}$ in place of $T_{i,a}$.

Lemma 2.2. *Let $W = (W, N, A, \triangleright, \mathbf{m})$ be the Coxeter groupoid associated with an element $(R, N, A, \triangleright)$ of $\mathcal{R}$ (see Definition 1.4). Assume that A is finite. Then there exists a map $f : W \to H_q(W)$ such that*

$$f(0) = 0, \ f(e_a) = E_a, \tag{26}$$

$$f(s_{i,a} w) = T_{i,a} f(w) \qquad \text{if } w \in W \setminus \{0\} \text{ and } \ell(s_{i,a} w) = 1 + \ell(w). \tag{27}$$

Further, as a $\mathbb{C}$-linear space, $H_q(W)$ is spanned by $f(W \setminus \{0\})$. In particular, if W is finite, then

$$\dim H_q(W) \leq |W| - 1. \tag{28}$$

Proof. Let $\widetilde{W}$ be the semigroup introduced in Definition 1.6 for W. It is easy to show that there exists a unique semigroup homomorphism $\widetilde{f} : \widetilde{W} \to H_q(W)$ such that $\widetilde{f}(0) = 0$, $\widetilde{f}(\tilde{e}_a) = E_a$ and $\widetilde{f}(\tilde{s}_{i,a}) = T_{i,a}$. By Theorem 1.7, there exists a unique map $f : W \to H_q(W)$ such that $f(0) = 0$ and $f(w) = \widetilde{f}(\tilde{s}_{i_1} \cdots \tilde{s}_{i_{m-1}} \tilde{s}_{i_m,a})$ if $w \in W \setminus \{0\}$, $\ell(w) = m$ and $w = s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a}$. Then f satisfies (26)-(27), as desired.

We show

$$\forall w \in W, \forall i \in N, \forall a \in A, T_{i,a} f(w) \in \mathbb{C} f(s_{i,a} w) + \mathbb{C} f(w). \tag{29}$$

If $s_{i,a}w = 0$, then clearly $T_{i,a}f(w) = 0$ holds. If $w \neq 0$, $s_{i,a}w \neq 0$ and $\ell(s_{i,a}w) = 1 + \ell(w)$, then (29) follows from (27). Assume that $w \neq 0$, $s_{i,a}w \neq 0$ and $\ell(s_{i,a}w) \neq 1+\ell(w)$. Then by (8) and Proposition 1.9, we have $\ell(s_{i,a}w) = \ell(w)-1$, so $f(w) = T_{i,i\triangleright a}f(s_{i,a}w)$. Since $T_{i,a}f(w) = T_{i,a}T_{i,i\triangleright a}f(s_{i,a}w)$, we have $T_{i,a}f(w) = f(s_{i,a}w)$ if $i \triangleright a \neq a$, and $T_{i,a}f(w) = (q-1)f(w) + qf(s_{i,a}w)$ otherwise. Hence we have (29), as desired.

It is clear from (29) that the rest of the statement follows. □

Notation 2.3. Let $r \in \mathbb{N}$. Let $V_0^{(r)}$ be the r-dimensional $\mathbb{R}$-linear space with a basis $\{\varepsilon_i | 1 \leq i \leq r\}$. Let $V_0^{(r),\prime}$ be the subspace of $V_0^{(r)}$ formed by the elements $\sum_{i=1}^{r} x_i\varepsilon_i$ with $x_i \in \mathbb{R}$ and $\sum_{i=1}^{r} x_i = 0$, so $\dim V_0^{(r),\prime} = r-1$. For a non-zero element $x = \sum_{i=1}^{|N|} x_i\varepsilon_i$ of $V_0^{(r)}$ with $x_i \in \mathbb{R}$, define $\widetilde{\sigma}_x \in \mathrm{GL}(V_0^{(r)})$ by $\widetilde{\sigma}_x(\varepsilon_j) = \varepsilon_j - 2x_j(\sum_{i=1}^{|N|} x_i^2)^{-1}x$, that is, $\widetilde{\sigma}_x$ is the reflection of $V_0^{(r)}$ with respect to the hyperplane of $V_0^{(r)}$ orthogonal to x. Note that if $x \in V_0^{(r),\prime}$, then $\widetilde{\sigma}_x(V_0^{(r),\prime}) = V_0^{(r),\prime}$.

2.4 Basic of Iwahori-Hecke algebras

For the basic facts about the Iwahori-Hecke algebras, we refer to [GU]. Let $W = (W, N, A, \triangleright, \mathbf{m})$ be the groupoid introdued in (1). In this subsection we always assume that

$$|A| = 1 \text{ and } N \text{ and } W \text{ are finite.} \tag{30}$$

Let $a \in A$, so $A = \{a\}$. Then $W \setminus \{0\}$ is nothing but the Coxeter group associated with the Coxeter system $(W \setminus \{0\}, \{s_{i,a}|, i \in N\})$. In this case, we also denote $H_q(W)$ and $T_{i,a}$ by $H_q(W \setminus \{0\})$ and T_i respectively. That is, $H_q(W \setminus \{0\})$ is the $\mathbb{C}$-algebra (with 1) generated by T_i $(i \in N)$ and defined by the relations

$$(T_i - q)(T_i + 1) = 0, \tag{31}$$

$$\begin{array}{lll} T_iT_j\cdots T_jT_i = T_jT_i\cdots T_iT_j \ (m_{i,j;a} \text{ factors}) & & \text{if } m_{i,j;a} \text{ is odd,} \\ T_jT_i\cdots T_jT_i = T_iT_j\cdots T_iT_j \ (m_{i,j;a} \text{ factors}) & & \text{if } m_{i,j;a} \text{ is even.} \end{array} \tag{32}$$

It is well-known that $\dim H_q(W\setminus\{0\}) = |W\setminus\{0\}|$. In this paper we fix a complete set of non-equivalent irreducible representations of $H_q(W \setminus \{0\})$ by

$$\{\rho_{H_q(W\setminus\{0\}),\lambda} : H_q(W \setminus \{0\}) \to \mathrm{End}_{\mathbb{C}}(V_{H_q(W\setminus\{0\}),\lambda}) | \lambda \in \Lambda_{H_q(W\setminus\{0\})}\}, \tag{33}$$

where $\Lambda_{H_q(W\setminus\{0\})}$ is an index set. Define the polynomial $P_{W\setminus\{0\}}(q)$ in q by

$$P_{W\setminus\{0\}}(q) := \sum_{w\in W\setminus\{0\}} q^{\ell(w)}. \tag{34}$$

This is called the *Poincaré polynomial* of $W \setminus \{0\}$.

It is well-known [GU] (see also [CR, (25.22) and (27.4)]) that for $q \in \mathbb{C} \setminus \{0\}$, the following three conditions are equivalent.

(i) $P_{W\setminus\{0\}}(q) \neq 0$ holds.
(ii) $H_q(W \setminus \{0\})$ is a semisimle algebra.
(iii) The map

$$\bigoplus_{\lambda\in\Lambda_{H_q(W\setminus\{0\})}} \rho_{H_q(W\setminus\{0\}),\lambda} : H_q(W \setminus \{0\}) \to \bigoplus_{\lambda\in\Lambda_{H_q(W\setminus\{0\})}} \mathrm{End}_{\mathbb{C}}(V_{H_q(W\setminus\{0\}),\lambda}) \tag{35}$$

defined by $X \mapsto \oplus_{\lambda\in\Lambda_{H_q(W\setminus\{0\})}}\rho_{H_q(W\setminus\{0\}),\lambda}(X)$ is a $\mathbb{C}$-algebra isomorphism.

In particular,

$$q \cdot P_{W\setminus\{0\}}(q) \neq 0 \Longrightarrow \dim H_q(W \setminus \{0\}) = \sum_{\lambda\in\Lambda_{H_q(W\setminus\{0\})}} (\dim V_{H_q(W\setminus\{0\}),\lambda})^2. \tag{36}$$

Assume that $N = \{1, 2, \dots, n\}$ and $m_{i,i+1;a} = 3$ and $m_{i,j;a} = 2$ ($|j - i| \geq 2$). Then W is the Coxeter groupoid associated with $(R, N, A, \triangleright) \in \mathcal{R}$ such that $V_0 = V_0^{(n+1),\prime}$, $R_a^+ = \{\varepsilon_i - \varepsilon_j | 1 \leq i < j \leq n + 1\}$, $\alpha_{i,a} = \varepsilon_i - \varepsilon_{i+1}$ and $\sigma_{i,a} = \widetilde{\sigma}_{\alpha_{i,a}}$. As a group, $W \setminus \{0\}$ is isomorphic to the symmetric group S_{n+1}, so we also denote $W \setminus \{0\}$ by S_{n+1} by abuse of notation. Note that $\dim H_q(S_{n+1}) = (n + 1)!$.

Assume that $N = \{1, 2, \dots, n\}$ and $m_{i,i+1;a} = 3$ ($1 \leq i \leq n - 3$), $m_{n-1,n;a} = 4$ and $m_{i,j;a} = 2$ ($|j - i| \geq 2$). Then W is the Coxeter groupoid associated with $(R, N, A, \triangleright) \in \mathcal{R}$ such that $V_0 = V_0^{(n)}$, $R_a^+ = \{\varepsilon_i - \varepsilon_j, \varepsilon_i + \varepsilon_j | 1 \leq i < j \leq n\} \cup \{\varepsilon_i | 1 \leq i \leq n\}$, $\alpha_{i,a} = \varepsilon_i - \varepsilon_{i+1}$ ($1 \leq i \leq n - 1$), $\alpha_{n,a} = \varepsilon_n$ and $\sigma_{i,a} = \widetilde{\sigma}_{\alpha_{i,a}}$. We also denote $W \setminus \{0\}$ by $W(B_n)$ and $W(C_n)$. Note that $\dim H_q(W(B_n)) = 2^n n!$.

Assume that $N = \{1, 2, \dots, n\}$ and $m_{i,i+1;a} = 3$ ($1 \leq i \leq n - 2$), $m_{n-1,n;a} = 2$, $m_{n-2,n;a} = 3$ and $m_{i,j;a} = 2$ ($|j - i| \geq 2$ and $1 \leq i \leq n - 3$). We also denote $W \setminus \{0\}$ by $W(D_n)$. Note that $\dim H_q(W(D_n)) = 2^{n-1}n!$. Then W is the Coxeter groupoid associated with $(R, N, A, \triangleright) \in \mathcal{R}$ such that $V_0 = V_0^{(n)}$, $R_a^+ = \{\varepsilon_i - \varepsilon_j, \varepsilon_i + \varepsilon_j | 1 \leq i < j \leq n\}$, $\alpha_{i,a} = \varepsilon_i - \varepsilon_{i+1}$ ($1 \leq i \leq n - 1$), $\alpha_{n,a} = \varepsilon_{n-1} + \varepsilon_n$ and $\sigma_{i,a} = \widetilde{\sigma}_{\alpha_{i,a}}$.

It is well-known (cf. [C, Theoerem 10.2.3 and Proposition 10.2.5]) that

$$P_{S_{n+1}}(q) = \prod_{r=1}^{n} \frac{q^{r+1} - 1}{q - 1}, \tag{37}$$

$$P_{W(B_n)}(q) = \prod_{r=1}^{n} \frac{q^{2r} - 1}{q - 1}, \tag{38}$$

$$P_{W(D_n)}(q) = \frac{q^n - 1}{q - 1} \prod_{r=1}^{n-1} \frac{q^{2r} - 1}{q - 1}. \tag{39}$$

2.5 Iwahori-Hecke type algebra $H_q(A(m,n))$ associated with the Lie superalgebra $A(m,n)$

Let

$$\triangleright : S_{m+n+2} \times \mathcal{D}_{m+1|n+1} \to \mathcal{D}_{m+1|n+1}$$

denote the usual (left) action of the symmetric group S_{m+n+2} on $\mathcal{D}_{m+1|n+1}$ by permutations, that is, for $\sigma \in S_{m+n+2}$,

$$\sigma \triangleright (p_1, \ldots, p_{m+n+2}) = (p_{\sigma^{-1}(1)}, \ldots, p_{\sigma^{-1}(m+n+2)}).$$

Let $\sigma_i := (i, i+1) \in S_{m+n+2}$. Let W be the Coxeter groupoid associated with $(R, N, A, \triangleright) \in \mathcal{R}$ such that $N = \{1, 2, \ldots, m+n+1\}$, $A = \mathcal{D}_{m+1|n+1}$, $i \triangleright d = \sigma_i \triangleright d$, $V_0 = V_0^{(m+n+2),\prime}$, $R_d^+ = \{\varepsilon_i - \varepsilon_j | 1 \leq i < j \leq m+n+1\}$, $\alpha_{i,d} = \varepsilon_i - \varepsilon_{i+1}$ and $\sigma_{i,d} = \widetilde{\sigma}_{\alpha_{i,d}}$. Denote $H_q(W)$ by $H_q(A(m,n))$. Then $H_q(A(m,n))$ is the $\mathbb{C}$-algebra (with 1) generated by

$$\{E_d \,|\, d \in \mathcal{D}_{m+1|n+1}\} \cup \{T_{i,d} \,|\, 1 \leq i \leq m+n+1,\, d \in \mathcal{D}_{m+1|n+1}\} \tag{40}$$

and defined by the relations (18)-(23) and the relations

$$T_{i,\sigma_j\sigma_i \triangleright d} T_{j,\sigma_i \triangleright d} T_{i,d} = T_{j,\sigma_i\sigma_j \triangleright d} T_{i,\sigma_j \triangleright d} T_{j,d} \qquad \text{if } |i-j| = 1, \tag{41}$$

$$T_{i,\sigma_j \triangleright d} T_{j,d} = T_{j,\sigma_i \triangleright d} T_{i,d} \qquad \text{if } |i-j| \geq 2. \tag{42}$$

Define d_e, $d_o \in \mathcal{D}_{m+1|n+1}$ by

$$d_e := (\overbrace{0, \ldots, 0}^{m+1}, \overbrace{1, \ldots, 1}^{n+1}),\ d_o := (\overbrace{1, \ldots, 1}^{n+1}, \overbrace{0, \ldots, 0}^{m+1}). \tag{43}$$

For $d = (p_1, \ldots, p_{m+n+2}) \in \mathcal{D}_{m+1|n+1}$, define the two elements

$$\tau_{+.d}, \tau_{-.d} \in S_{m+n+2} \tag{44}$$

by

$$p_{\tau_{\pm.d}(i)} = \frac{1 \mp 1}{2} \quad \text{and} \quad \tau_{\pm.d}(i) \leq \tau_{\pm.d}(j) \qquad \text{if } 1 \leq i \leq j \leq m+1, \tag{45}$$

$$p_{\tau_{\pm.d}(i)} = \frac{1 \pm 1}{2} \quad \text{and} \quad \tau_{\pm.d}(i) \leq \tau_{\pm.d}(j) \qquad \text{if } m+2 \leq i \leq j \leq m+n+2. \tag{46}$$

Then $\tau_{+,d}$ (resp. $\tau_{-,d}$) is the minimal length element among the elements $\sigma \in S_{m+n+2}$ satisfying the condition that for any i, i-th component of d_e (resp. d_o) is the same as $\sigma(i)$-th component $p_{\sigma(i)}$ of d.

Example 2.4. Assume that $m = n = 1$. Then $\mathcal{D}_{2|2} = \{d_e = (0,0,1,1), d_1 = (0,1,0,1), d_2 = (1,0,0,1), d_3 = (0,1,1,0), d_4 = (1,0,1,0), d_o = (1,1,0,0)\}$. Then $\tau_{+.d_e} = \begin{bmatrix}1234\\1234\end{bmatrix}$, $\tau_{-.d_e} = \begin{bmatrix}1234\\3412\end{bmatrix}$, $\tau_{+.d_1} = \begin{bmatrix}1234\\1324\end{bmatrix}$, $\tau_{-.d_1} = \begin{bmatrix}1234\\2413\end{bmatrix}$, $\tau_{+.d_2} = \begin{bmatrix}1234\\2314\end{bmatrix}$, $\tau_{-.d_2} = \begin{bmatrix}1234\\1423\end{bmatrix}$, $\tau_{+.d_3} = \begin{bmatrix}1234\\1423\end{bmatrix}$, $\tau_{-.d_3} = \begin{bmatrix}1234\\2314\end{bmatrix}$, $\tau_{+.d_4} = \begin{bmatrix}1234\\2413\end{bmatrix}$, $\tau_{-.d_4} = \begin{bmatrix}1234\\1324\end{bmatrix}$, $\tau_{+.d_o} = \begin{bmatrix}1234\\3412\end{bmatrix}$, $\tau_{-.d_o} = \begin{bmatrix}1234\\1234\end{bmatrix}$. See also Figure 2.



Figure 2: Dynkin diagrams of the Lie superalgebra $A(1,1)$

Now we consider $|W|$. Recall ρ and d_e from Theorem 1.5 and (44) respectively. It is easy to see that $P_{d_e}\rho(e_{d_e}We_{d_e})\iota_{d_e} \subset \{(\sum_{i=1}^{m+n+2}\mathbf{E}_{\sigma(i)i})_{|V_0^{(m+n+2),\prime}}|\sigma \in S_{m+n+2}, \sigma(\{1,\ldots,m+1\}) = \{1,\ldots,m+1\}\}$. Hence $|e_{d_e}We_{d_e}| \leq (m+1)!(n+1)!$ by Theorem 1.5, so $|W \setminus \{0\}| = |\mathcal{D}_{m+1|n+1}|^2|e_{d_e}We_{d_e}| \leq \frac{((m+n+2)!)^2}{(m+1)!(n+1)!}$. Hence by (28), we conclude

$$\dim H_q(A(m,n)) \leq \frac{((m+n+2)!)^2}{(m+1)!(n+1)!}. \tag{47}$$

Proposition 2.5. *Let V and W be finite dimensional $\mathbb{C}$-linear spaces, and let $\mathbf{l} : H_q(S_{m+1}) \to \mathrm{End}_{\mathbb{C}}(V)$ and $\mathbf{r} : H_q(S_{n+1}) \to \mathrm{End}_{\mathbb{C}}(W)$ be $\mathbb{C}$-algebra homomorphisms, i.e., representations. Let $\mathbf{l}\otimes\mathbf{r} : H_q(S_{m+1}) \otimes H_q(S_{n+1}) \to \mathrm{End}_{\mathbb{C}}(V\otimes W)$ denote the tensor representation of $\mathbf{l}$ and $\mathbf{r}$ in the ordinary sense. Let $C_{V\otimes W;d}$ be copies of the $\mathbb{C}$-linear space $V\otimes W$, indexed by $d \in \mathcal{D}_{m+1|n+1}$. Let $C_{V\otimes W} := \oplus_{d\in\mathcal{D}_{m+1|n+1}} C_{V\otimes W;d}$. Let $P_d : C_{V\otimes W} \to C_{V\otimes W;d}$ and $\iota_d : C_{V\otimes W;d} \to C_{V\otimes W}$ denote the canonical projection and the canonical inclusion map respectively. Then there exists a unigue $\mathbb{C}$-algebra homomorphism $\mathbf{l}\boxtimes^{A(m,n)}\mathbf{r} : H_q(A(m,n)) \to \mathrm{End}_{\mathbb{C}}(C_{V\otimes W})$ satisfying the following conditions:*

(i) *For each $d \in \mathcal{D}_{m+1|n+1}$, one has $(\mathbf{l}\boxtimes^{A(m,n)}\mathbf{r})(E_d) = \iota_d \circ P_d$,*

(ii) *For each $i \in \{1,\ldots,m+n+1\}$ and each $d = (p_1,\ldots,p_{m+n+2}) \in \mathcal{D}_{m+1|n+1}$, one has*

$$(\mathbf{l}\boxtimes^{A(m,n)}\mathbf{r})(T_{i,d}) = \begin{cases} P_{\sigma_i\triangleright d}\circ\iota_d & \text{if } p_i \neq p_{i+1}, \\ \iota_d\circ(\mathbf{l}(T_{\tau_{+.d}^{-1}(i)})\otimes\mathrm{id}_W)\circ P_d & \text{if } p_i = p_{i+1} = 0, \\ \iota_d\circ(\mathrm{id}_V\otimes\mathbf{r}(T_{\tau_{-.d}^{-1}(i)}))\circ P_d & \text{if } p_i = p_{i+1} = 1. \end{cases} \tag{48}$$

Proof. This can be checked directly. Refer to Figure 3. We explain by using an example. Denote $(\mathbf{l}\boxtimes^{A(m,n)}\mathbf{r})(T_{i',d'})$ by $S_{i',d'}$ for any d' and i'. Let


p_i p_{i+1} p_{i+2} $\in \{0,1\}$
d
$i \triangleright d$
$(i+1) \triangleright i \triangleright d$
$i \triangleright (i+1) \triangleright i \triangleright d$
p_{i+2} p_{i+1} p_i
p_i p_{i+1} p_{i+2} $\in \{0,1\}$
d
$(i+1) \triangleright d$
$i \triangleright (i+1) \triangleright d$
$(i+1) \triangleright i \triangleright (i+1) \triangleright d$
p_{i+2} p_{i+1} p_i

Figure 3: Braid relation

$d = (p_1, \dots, p_{m+n+2}) \in \mathcal{D}_{m+1|n+1}$ and $i \in \{1, \dots, m+n\}$ and assume $p_i = p_{i+1} = 0$ and $p_{i+2} = 1$. Let $d_1 := i \triangleright d (= \sigma_i \triangleright d)$, $d_2 := (i+1) \triangleright d_1$, $d_3 := i \triangleright d_2$, $d_4 := (i+1) \triangleright d$, $d_5 := i \triangleright d_4$ and $d_6 := (i+1) \triangleright d_4$. Then

$$d = d_1 = (p_1, \dots, p_{i-1}, 0, 0, 1, p_{i+2}, \dots, p_{m+n+2}), \tag{49}$$

$$d_2 = d_4 = (p_1, \dots, p_{i-1}, 0, 1, 0, p_{i+2}, \dots, p_{m+n+2}), \tag{50}$$

$$d_3 = d_5 = d_6 = (p_1, \dots, p_{i-1}, 1, 0, 0, p_{i+2}, \dots, p_{m+n+2}). \tag{51}$$

Note that $\tau_{+.d_5} = \sigma_i \sigma_{i+1} \tau_{+.d}$. Hence $\tau_{+.d_5}^{-1}(i+1) = \tau_{+.d}^{-1}(i)$. Then we have $S_{i,d} = \iota_d \circ (\mathbf{l}(T_{\tau_{+.d}^{-1}(i)}) \otimes \mathrm{id}_W) \circ P_d$, $S_{i+1,d_1} = \iota_{d_2} \circ P_d$, $S_{i,d_2} = \iota_{d_3} \circ P_{d_2}$, $S_{i+1,d} = \iota_{d_2} \circ P_d$, $S_{i,d_4} = \iota_{d_3} \circ P_{d_2}$ and $S_{i+1,d_5} = \iota_{d_3} \circ (\mathbf{l}(T_{\tau_{+.d}^{-1}(i)}) \otimes \mathrm{id}_W) \circ P_{d_3}$. Hence we have $S_{i,d_2} S_{i+1,d_1} S_{i,d} = S_{i+1,d_5} S_{i,d_4} S_{i+1,d} = \iota_{d_3} \circ (\mathbf{l}(T_{\tau_{+.d}^{-1}(i)}) \otimes \mathrm{id}_W) \circ P_d$, as desired. □

For $\lambda \in \Lambda_{H_q(S_{m+1})}$ and $\mu \in \Lambda_{H_q(S_{n+1})}$, we denote $\rho_{H_q(S_{m+1}),\lambda} \boxtimes^{A(m,n)} \rho_{H_q(S_{n+1}),\mu}$ by $\rho^{A(m,n)}_{q;\lambda,\mu}$ and we denote $C_{V\otimes W}$, P_d, ι_d for $V = V_{H_q(S_{m+1}),\lambda}$ and $W = V_{H_q(S_{n+1}),\mu}$ by $C^{A(m,n)}_{q;\lambda,\mu}$, $P^{\lambda,\mu}_d$, $\iota^{\lambda,\mu}_d$ respectively.

Theorem 2.6. *Let $q \in \mathbb{C}$ and assume that*

$$qP_{S_{m+1}}(q)P_{S_{n+1}}(q) \neq 0. \tag{52}$$

Then the $\mathbb{C}$-algebra homomorphism

$$\begin{aligned}(53)\qquad & \bigoplus_{(\lambda,\mu)\in\Lambda_{H_q(S_{m+1})}\times\Lambda_{H_q(S_{n+1})}} \rho^{A(m,n)}_{q;\lambda,\mu} : \\ & H_q(A(m,n)) \to \bigoplus_{(\lambda,\mu)\in\Lambda_{H_q(S_{m+1})}\times\Lambda_{H_q(S_{n+1})}} \mathrm{End}_{\mathbb{C}}(C^{A(m,n)}_{q;\lambda,\mu})\end{aligned}$$

is an isomorphism. Further we have

$$\dim H_q(A(m,n)) = \frac{((m+n+2)!)^2}{(m+1)!(n+1)!}. \tag{54}$$

Moreover $H_q(A(m,n))$ is a semisimple $\mathbb{C}$-algebra and a complete set of non-equivalent irreducible representations of $H_q(A(m,n))$ is given by $\{\rho^{A(m,n)}_{q;\lambda,\mu} | (\lambda,\mu) \in \Lambda_{H_q(S_{m+1})} \times \Lambda_{H_q(S_{n+1})}\}$.

Proof. Define the $\mathbb{C}$-algebra homomorphism $f_1 : H_q(S_{m+1}) \otimes H_q(S_{n+1}) \to H_q(A(m,n))$ by $f_1(T_i \otimes 1) = T_{i,e}$ and $f_1(1 \otimes T_j) = T_{m+1+j,e}$. Let $R_{\lambda,\mu} := (\iota^{\lambda,\mu}_{d_e} \circ P^{\lambda,\mu}_{d_e})\mathrm{End}_{\mathbb{C}}(C^{A(m,n)}_{q;\lambda,\mu})(\iota^{\lambda,\mu}_{d_e} \circ P^{\lambda,\mu}_{d_e})$. Let f_2 denote the homomorphism of (53). It follows from (52) that $H_q(S_{m+1}) \otimes H_q(S_{n+1})$ is a semisimple $\mathbb{C}$-algebra. This implies

$$\mathrm{Im}(f_2 \circ f_1) = \bigoplus_{(\lambda,\mu)\in\Lambda_{H_q(S_{m+1})}\times\Lambda_{H_q(S_{n+1})}} R_{\lambda,\mu}.$$

On the other hand, we have

$$\mathrm{End}_{\mathbb{C}}(C^{A(m,n)}_{q;\lambda,\mu}) = \bigoplus_{d_1,d_2\in\mathcal{D}_{m+1|n+1}} (\iota^{\lambda,\mu}_{d_1} \circ P^{\lambda,\mu}_{d_e}) R_{\lambda,\mu} (\iota^{\lambda,\mu}_{d_e} \circ P^{\lambda,\mu}_{d_2}).$$

Hence by (48) we can easily see that f_2 is surjective. In particular, we have

$$\begin{aligned}
&\dim H_q(A(m,n)) \\
&\geq \sum_{(\lambda,\mu)\in\Lambda_{H_q(S_{m+1})}\times\Lambda_{H_q(S_{n+1})}} |\mathcal{D}_{m+1|n+1}|^2 \dim R_{\lambda,\mu} \\
&= |\mathcal{D}_{m+1|n+1}|^2 \sum_{(\lambda,\mu)\in\Lambda_{H_q(S_{m+1})}\times\Lambda_{H_q(S_{n+1})}} \dim R_{\lambda,\mu} \\
&= (\frac{(m+n+2)!}{(m+1)!(n+1)!})^2 \\
&\qquad \sum_{(\lambda,\mu)\in\Lambda_{H_q(S_{m+1})}\times\Lambda_{H_q(S_{n+1})}} (\dim V_{H_q(S_{m+1}),\lambda})^2 (\dim V_{H_q(S_{n+1}),\mu})^2 \\
&= (\frac{(m+n+2)!}{(m+1)!(n+1)!})^2 \dim H_q(S_{m+1}) \dim H_q(S_{n+1}) \\
&= \frac{((m+n+2)!)^2}{(m+1)!(n+1)!}.
\end{aligned}$$



Figure 4: Dynkin diagrams of the Lie superalgebra $B(1,2)$

Hence by (47), we have (54). Hence f_2 is an isomorphism. Then the rest of the statement follows from well-known facts concerning semisimple algebras (cf. [CR, (25.22) and (27.4)]). $\square$

2.6 Iwahori-Hecke type algebra associated with the Lie superalgebra $\mathfrak{osp}(2m+1|2n)$

Let $m \in \mathbb{N} \cup \{0\}$ and $n \in \mathbb{N}$. Ler $\ell := m+n$. For $1 \leq i \leq \ell$, define $\hat{\sigma}_i \in S_\ell$ by $\hat{\sigma}_i := \sigma_i$ $(1 \leq i \leq \ell - 1)$. and $\hat{\sigma}_\ell := \mathrm{id}$.

Let W be the Coxeter groupoid associated with $(R, N, A, \triangleright) \in \mathcal{R}$ such that $N = \{1, 2, \ldots, \ell\}$, $A = \mathcal{D}_{m|n}$, $i \triangleright d = \hat{\sigma}_i \triangleright d$, $V_0 = V_0^{(m+n)}$, $R_d^+ = \{\varepsilon_i - \varepsilon_j, \varepsilon_i - \varepsilon_j | 1 \leq i < j \leq \ell\} \cup \{\varepsilon_i | 1 \leq i \leq \ell\}$, $\alpha_{i,d} = \varepsilon_i - \varepsilon_{i+1}$ $(1 \leq i \leq \ell - 1)$, $\alpha_{\ell,d} = \varepsilon_\ell$ and $\sigma_{i,d} = \widetilde{\sigma}_{\alpha_{i,d}}$. Denote $H_q(W)$ by $H_q(B(m,n))$. Then $H_q(B(m,n))$ is the $\mathbb{C}$-algebra (with 1) generated by

$$\{E_d \,|\, d \in \mathcal{D}_{m|n}\} \cup \{T_{i,d} \,|\, 1 \leq i \leq \ell,\, d \in \mathcal{D}_{m|n}\} \tag{55}$$

and defined by the relations (18)-(23) and the relations

$$T_{\ell-1,\hat{\sigma}_{\ell-1}\triangleright d} T_{\ell,\hat{\sigma}_{\ell-1}\triangleright d} T_{\ell-1,d} T_{\ell,d} = T_{\ell,d} T_{\ell-1,\hat{\sigma}_{\ell-1}\triangleright d} T_{\ell,\hat{\sigma}_{\ell-1}\triangleright d} T_{\ell-1,d} \tag{56}$$

$$T_{i,\hat{\sigma}_{i+1}\hat{\sigma}_i\triangleright d} T_{i+1,\hat{\sigma}_i\triangleright d} T_{i,d} = T_{i+1,\hat{\sigma}_i\hat{\sigma}_{i+1}\triangleright d} T_{i,\hat{\sigma}_{i+1}\triangleright d} T_{i+1,d} \qquad \text{if } 1 \leq i \leq \ell - 1, \tag{57}$$

$$T_{i,\hat{\sigma}_j\triangleright d} T_{j,d} = T_{j,\hat{\sigma}_i\triangleright d} T_{i,d} \qquad \text{if } |i - j| \geq 2. \tag{58}$$

Recall ρ and $d_e \in \mathcal{D}_{m|n}$ from Theorem 1.5 and (43) respectively. Then $P_{d_e}\rho(e_{d_e} W e_{d_e})\iota_{d_e} \subset \{(\sum_{i=1}^{m+n} z_i \mathbf{E}_{\sigma(i)i}) | \sigma \in S_{m+n}, z_i \in \{-1, 1\}, \sigma(\{1, \ldots, m\}) =$

$\{1,\ldots,m\}\}$. Hence $|e_{d_e}We_{d_e}| \leq 2^{m+n}m!n!$ by Theorem 1.5, so $|W \setminus \{0\}| = |\mathcal{D}_{m|n}|^2|e_{d_e}We_{d_e}| \leq \frac{2^{m+n}((m+n)!)^2}{m!n!}$. Hence by (28), we conclude

$$\dim H_q(B(m,n)) \leq \frac{2^{m+n}((m+n)!)^2}{m!n!}. \tag{59}$$

Proposition 2.7. *Let* $V_{\mathbf{l}}$ *and* $V_{\mathbf{r}}$ *be finite dimensional* $\mathbb{C}$*-linear spaces, and let* $\mathbf{l} : H_q(W(B_m)) \to \mathrm{End}_{\mathbb{C}}(V_{\mathbf{l}})$ *and* $\mathbf{r} : H_q(W(B_n)) \to \mathrm{End}_{\mathbb{C}}(V_{\mathbf{r}})$ *be* $\mathbb{C}$*-algebra homomorphisms, i.e., representations. Let* $\mathbf{l} \otimes \mathbf{r} : H_q(W(B_m)) \otimes H_q(W(B_n)) \to \mathrm{End}_{\mathbb{C}}(V_{\mathbf{l}} \otimes V_{\mathbf{r}})$ *denote the tensor representation of* $\mathbf{l}$ *and* $\mathbf{r}$ *in the ordinary sense. Let* $C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}};d}$ *be copies of the* $\mathbb{C}$*-linear space* $V_{\mathbf{l}} \otimes V_{\mathbf{r}}$*, indexed by* $d \in \mathcal{D}_{m|n}$*. Let* $C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}}} := \oplus_{d \in \mathcal{D}_{m|n}} C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}};d}$*. Let* $P_d : C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}}} \to C_{V \otimes W;d}$ *and* $\iota_d : C_{V \otimes W;d} \to C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}}}$ *denote the canonical projection and the canonical inclusion map respectively. Then there exists a unique* $\mathbb{C}$*-algebra homomorphism* $\mathbf{l} \boxtimes \mathbf{r} = \mathbf{l} \boxtimes^{B(m,n)} \mathbf{r} : H_q(B(m,n)) \to \mathrm{End}_{\mathbb{C}}(C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}}})$ *satisfying the following conditions:*

(i) *For each* $d \in \mathcal{D}_{m|n}$*, one has* $(\mathbf{l} \boxtimes \mathbf{r})(E_d) = \iota_d \circ P_d$*,*

(ii) *For each* $i \in \{1,\ldots,\ell = m+n\}$ *and each* $d = (p_1,\ldots,p_\ell) \in \mathcal{D}_{m|n}$*, one has*

$$(\mathbf{l} \boxtimes \mathbf{r})(T_{i,d}) = \begin{cases} P_{\hat{\sigma}_i \triangleright d} \circ \iota_d & \text{if } 1 \leq i \leq \ell - 1 \text{ and } p_i \neq p_{i+1}, \\ \iota_d \circ (\mathbf{l}(T_{\tau_{+.d}^{-1}(i)}) \otimes \mathrm{id}_{V_{\mathbf{r}}}) \circ P_d & \text{if } 1 \leq i \leq \ell - 1 \text{ and } p_i = p_{i+1} = 0, \\ \iota_d \circ (\mathrm{id}_{V_{\mathbf{l}}} \otimes \mathbf{r}(T_{\tau_{-.d}^{-1}(i)})) \circ P_d & \text{if } 1 \leq i \leq \ell - 1 \text{ and } p_i = p_{i+1} = 1, \\ \iota_d \circ (\mathbf{l}(T_m) \otimes \mathrm{id}_{V_{\mathbf{r}}}) \circ P_d & \text{if } i = \ell \text{ and } p_\ell = 0, \\ \iota_d \circ (\mathrm{id}_{V_{\mathbf{l}}} \otimes \mathbf{r}(T_n)) \circ P_d & \text{if } i = \ell \text{ and } p_\ell = 1, \end{cases} \tag{60}$$

where $\tau_{\pm.d}$ *are the ones of* (44).

Proof. We can check out this directly in a way similar to that for Proof of Proposition 2.5. □

For $\lambda \in \Lambda_{H_q(W(B_m))}$ and $\mu \in \Lambda_{H_q(W(B_n))}$, we denote $\rho_{H_q(W(B_m)),\lambda} \boxtimes^{B(m,n)} \rho_{W(B_n)),\mu}$ by $\rho^{B(m,n)}_{q;\lambda,\mu}$ and we denote $C_{V \otimes W}$ for $V = V_{H_q(W(B_m)),\lambda}$ and $W = V_{H_q(W(B_n)),\mu}$ by $C^{B(m,n)}_{q;\lambda,\mu}$.

Theorem 2.8. *Let* $q \in \mathbb{C}$ *and assume that*

$$qP_{W(B_m)}(q)P_{W(B_n)}(q) \neq 0. \tag{61}$$

Then the $\mathbb{C}$-algebra homomorphism

$$\bigoplus_{(\lambda,\mu)\in\Lambda_{H_q(W(B_m))}\times\Lambda_{H_q(W(B_n))}} \rho^{B(m,n)}_{q;\lambda,\mu} : \quad H_q(B(m,n)) \to \bigoplus_{(\lambda,\mu)\in\Lambda_{H_q(W(B_m))}\times\Lambda_{H_q(W(B_n))}} \mathrm{End}_{\mathbb{C}}(C^{B(m,n)}_{q;\lambda,\mu}) \tag{62}$$

is an isomorphism. Further we have

$$\dim H_q(B(m,n)) = \frac{2^{m+n}((m+n)!)^2}{m!n!}. \tag{63}$$

Moreover $H_q(B(m,n))$ is a semisimple $\mathbb{C}$-algebra and a complete set of non-equivalent irreducible representations of $H_q(B(m,n))$ is given by $\{\rho^{B(m,n)}_{q;\lambda,\mu}|(\lambda,\mu)\in\Lambda_{H_q(W(B_m))}\times\Lambda_{H_q(W(B_n))}\}$.

Proof. Let $\mathbf{l} : H_q(W(B_m)) \to \mathrm{End}_{\mathbb{C}}(V_{\mathbf{l}})$ and $\mathbf{r} : H_q(W(B_n)) \to \mathrm{End}_{\mathbb{C}}(V_{\mathbf{r}})$ be irreducible representations. Further, let $\mathbf{l}\boxtimes\mathbf{r} : H_q(B(m,n)) \to \mathrm{End}_{\mathbb{C}}(C_{V_{\mathbf{l}}\otimes V_{\mathbf{r}}})$ be the representaion introduced in Proposition 2.7 for these $\mathbf{l}$ and $\mathbf{r}$. By (60), we can easily see that

$$\forall d', \forall d'' \in \mathcal{D}_{m|n}, \quad P_{d'}\circ\iota_{d''} \in \mathrm{Im}(\mathbf{l}\boxtimes\mathbf{r}). \tag{64}$$

Define the representation $f_1 : H_q(W(B_m))\otimes H_q(W(B_n)) \to \mathrm{End}_{\mathbb{C}}(C_{V_{\mathbf{l}}\otimes V_{\mathbf{r}};d_e})$ by $f_1(T_i\otimes 1) = (P_{d_e}\circ\iota_{d_o})((\mathbf{l}\boxtimes\mathbf{r})(T_{n+i,d_o}))(P_{d_o}\circ\iota_{d_e})$ and $f_1(1\otimes T_j) = (\mathbf{l}\boxtimes\mathbf{r})(T_{m+j,d_e})$. The condition (61) implies that f_1 is an irreducible representaion of $H_q(W(B_m))\otimes H_q(W(B_n))$. Moreover, using (64), we can easily see that $\mathbf{l}\boxtimes\mathbf{r}$ is an irreducible representation of $H_q(B(m,n))$.

By the above argument, together with (59), in the same way as that for Proof of Theorem 2.6, we can complete the proof of this theorem. □

2.7 Iwahori-Hecke type algebra associated with the Lie superalgebra $\mathfrak{osp}(2m|2n)$

Let m, $n\in\mathbb{N}$. Define the set $\mathcal{D}^{CD}_{m|n}$ by

$$\begin{aligned}\mathcal{D}^{CD}_{m|n} :=&\{d^D|d=(p_1,\ldots,p_{m+n})\in\mathcal{D}_{m|n},\, p_{m+n}=0\}\\ &\cup\{d^C_+, d^C_-|d=(p_1,\ldots,p_{m+n})\in\mathcal{D}_{m|n},\, p_{m+n}=1\},\end{aligned} \tag{65}$$

so that

$$|\mathcal{D}^{CD}_{m|n}| = \frac{(m+n-1)!}{(m-1)!n!} + 2\frac{(m+n-1)!}{m!(n-1)!} = \frac{(m+n-1)!(m+2n)}{m!n!}. \tag{66}$$

Let $\ell := m+n$ and $N = \{1, \dots, \ell\}$. Define the action $\triangleright$ of $F_2(N)$ on $\mathcal{D}^{CD}_{m|n}$ by

$$
i \triangleright a = \begin{cases} (\sigma_i \triangleright d)^D & \text{if } a = d^D,\ 1 \le i \le \ell-2 \text{ and } p_i \ne p_{i+1}, \\ (\sigma_i \triangleright d)^C_+ & \text{if } a = d^D,\ i = \ell-1 \text{ and } p_i \ne p_{i+1}, \\ (\sigma_{i-1} \triangleright d)^C_- & \text{if } a = d^D,\ i = \ell \text{ and } p_{i-1} \ne p_i, \\ (\sigma_i \triangleright d)^C_\pm & \text{if } a = d^C_\pm,\ 1 \le i \le \ell-2 \text{ and } p_i \ne p_{i+1}, \\ (\sigma_i \triangleright d)^D & \text{if } a = d^C_+,\ i = \ell-1 \text{ and } p_i \ne p_{i+1}, \\ (\sigma_{i-1} \triangleright d)^D & \text{if } a = d^C_-,\ i = \ell \text{ and } p_{i-1} \ne p_i, \\ a & \text{otherwise.} \end{cases} \tag{67}
$$

Now we define $R = (R, N, \mathcal{D}^{CD}_{m|n}, \triangleright) \in \mathcal{R}$ as follows. Let N be as above. Let $A = \mathcal{D}^{CD}_{m|n}$. Let $V_0 = V_0^{(\ell)}$. Let $a = d^D$, d^C_+ or $d^C_- \in \mathcal{D}^{CD}_{m|n}$ with $d = (p_1, \dots, p_{m+n}) \in \mathcal{D}_{m|n}$. Let R^+_a be the subset of V_0 formed by the elements $\varepsilon_i - \varepsilon_j$, $\varepsilon_i + \varepsilon_j$, $(1 \le i < j \le \ell)$ and $2\varepsilon_k$ $(1 \le k \le \ell$ and $p_k = 1)$. Define

$$
\alpha_{i,a} := \begin{cases} \varepsilon_i - \varepsilon_{i+1} & \text{if } a = d^D \text{ or } d^C_+ \text{ and } 1 \le i \le \ell-1, \\ \varepsilon_i - \varepsilon_{i+1} & \text{if } a = d^C_- \text{ and } 1 \le i \le \ell-2, \\ \varepsilon_{\ell-1} + \varepsilon_\ell & \text{if } a = d^D \text{ and } i = \ell, \\ 2\varepsilon_\ell & \text{if } a = d^C_+ \text{ and } i = \ell, \\ -2\varepsilon_\ell & \text{if } a = d^C_- \text{ and } i = \ell-1, \\ \varepsilon_{\ell-1} + \varepsilon_\ell & \text{if } a = d^C_- \text{ and } i = \ell. \end{cases} \tag{68}
$$

Define $\sigma_{i.a} := \widetilde{\sigma}_{\alpha_{i.a}}$. Let W be the Coxeter groupoid associated with R. Recall ρ and $d_e \in \mathcal{D}_{m|n}$ from Theorem 1.5 and (43) respectively. It is easy to show that

$$
\rho(e_{(d_e)^D} W e_{(d_e)^D}) = \\ \{\sum_{j=1}^{\ell} z_j \mathbf{E}_{\sigma(j)j} | \sigma \in S_\ell, z_j \in \{-1, 1\}, \prod_{j=n+1}^{\ell} z_j = 1, \sigma(\{1, \dots, n\}) = \{1, \dots, n\}\},
$$

so $|e_{(d_e)^D} W e_{(d_e)^D}| \le m!n!2^{\ell-1}$ by Theorem 1.5. Hence $|W \backslash \{0\}| \le |\mathcal{D}^{CD}_{m|n}|^2 m!n!2^{\ell-1}$. Denote $H_q(W)$ by $H_q(\mathfrak{osp}(2m|2n))$. By (28) and (66), we have

$$
\dim H_q(\mathfrak{osp}(2m|2n)) \le \frac{2^{m+n-1}((m+n-1)!(m+2n))^2}{m!n!}. \tag{69}
$$

Recall that $H_q(\mathfrak{osp}(2m|2n))$ is the $\mathbb{C}$-algebra (with 1) generated by

$$
\{E_a \,|\, a \in \mathcal{D}^{CD}_{m|n}\} \cup \{T_{i,a} \,|\, 1 \le i \le m+n,\ a \in \mathcal{D}^{CD}_{m|n}\} \tag{70}
$$



Figure 5: Dynkin diagrams of the Lie superalgebra $D(3,1)$

and defined by the relations (18)-(23) and the relations

(71) $(T_{i,a}T_{j,a})^2 = (T_{j,a}T_{i,a})^2$ if $a = d^C_{\pm}$, $p_{\ell-1} = p_\ell$ and $i = \ell - 1$, $j = \ell$,

(72) $T_{i,a}T_{j,a} = T_{j,a}T_{i,a}$ if $a = d^D$, $p_{\ell-1} = p_\ell$ and $i = \ell - 1$, $j = \ell$,

(73) $T_{i,ji\triangleright d}T_{j,i\triangleright d}T_{i,d} = T_{j,ij\triangleright d}T_{i,j\triangleright d}T_{j,d}$ if $p_{\ell-1} \neq p_\ell$ and $i = \ell - 1$, $j = \ell$,

(74) $T_{i,ji\triangleright d}T_{j,i\triangleright d}T_{i,d} = T_{j,ij\triangleright d}T_{i,j\triangleright d}T_{j,d}$ if $1 \leq i \leq \ell - 3$, $j = i + 1$,

(75) $T_{i,ji\triangleright d}T_{j,i\triangleright d}T_{i,d} = T_{j,ij\triangleright d}T_{i,j\triangleright d}T_{j,d}$ if $a = d^C_+$ and $i = \ell - 2$, $j = \ell - 1$,

(76) $T_{i,ji\triangleright d}T_{j,i\triangleright d}T_{i,d} = T_{j,ij\triangleright d}T_{i,j\triangleright d}T_{j,d}$ if $a = d^C_-$ and $i = \ell - 2$, $j = \ell$,

(77) $T_{i,ji\triangleright d}T_{j,i\triangleright d}T_{i,d} = T_{j,ij\triangleright d}T_{i,j\triangleright d}T_{j,d}$ if $a = d^D$ and $i = \ell - 2$, $j \in \{\ell - 1, \ell\}$,

(78) $T_{j,i\triangleright d}T_{i,d} = T_{i,j\triangleright d}T_{j,d}$ if $i < j$ holds and i, j are not the ones in (76) - (77).

Recall that $W(C_k) = W(B_k)$ and $H_q(W(C_k)) = H_q(W(B_k))$.

Proposition 2.9. *Let $V_{\mathbf{l}}$ and $V_{\mathbf{r}}$ be finite dimensional $\mathbb{C}$-linear spaces, and let $\mathbf{l} : H_q(W(D_m)) \to \mathrm{End}_{\mathbb{C}}(V_{\mathbf{l}})$ and $\mathbf{r} : H_q(W(C_n)) \to \mathrm{End}_{\mathbb{C}}(V_{\mathbf{r}})$ be $\mathbb{C}$-algebra homomorphisms, i.e., representations. Let $\mathbf{l} \otimes \mathbf{r} : H_q(W(D_m)) \otimes H_q(W(C_n)) \to \mathrm{End}_{\mathbb{C}}(V_{\mathbf{l}} \otimes V_{\mathbf{r}})$ denote the tensor representation of $\mathbf{l}$ and $\mathbf{r}$ in the ordinary sense. Let $C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}};a}$ be copies of the $\mathbb{C}$-linear space $V_{\mathbf{l}} \otimes V_{\mathbf{r}}$, indexed by $a \in \mathcal{D}^{CD}_{m|n}$. Let $C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}}} := \oplus_{d \in \mathcal{D}^{CD}_{m|n}} C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}};d}$. Let $P_a : C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}}} \to C_{V \otimes W;a}$ and $\iota_a : C_{V \otimes W;a} \to C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}}}$ denote the canonical projection and the canonical inclusion map respectively. Then there exists a unique $\mathbb{C}$-algebra homomorphism $\mathbf{l} \boxtimes \mathbf{r} = \mathbf{l} \boxtimes^{CD} \mathbf{r} : H_q(\mathfrak{osp}(2m|2n)) \to \mathrm{End}_{\mathbb{C}}(C_{V_{\mathbf{l}} \otimes V_{\mathbf{r}}})$ satisfying the following conditions:*

(i) *For each $a \in \mathcal{D}^{CD}_{m|n}$, one has $(\mathbf{l} \boxtimes \mathbf{r})(E_a) = \iota_a \circ P_a$.*

(ii) *For each $i \in \{1, \ldots, \ell = m+n\}$ and each $a \in \mathcal{D}^{CD}_{m|n}$ with $d = (p_1, \ldots, p_\ell) \in$*

$\mathcal{D}_{m|n}$ such that $a = d^C_+$, d^C_- or d^D, one has

$$(\mathbf{l} \boxtimes \mathbf{r})(T_{i,a}) = \begin{cases} P_{i \triangleright a} \circ \iota_a & \text{if } 1 \leq i \leq \ell \text{ and } i \triangleright a \neq a, \\ \iota_a \circ (\mathbf{l}(T_{\tau^{-1}_{+.d}(i)}) \otimes \mathrm{id}_{V_\mathbf{r}}) \circ P_a & \text{if } 1 \leq i \leq \ell - 1 \text{ and } p_i = p_{i+1} = 0, \\ \iota_a \circ (\mathbf{l}(T_m) \otimes \mathrm{id}_{V_\mathbf{r}}) \circ P_a & \text{if } i = \ell \text{ and } p_\ell = 0, \\ \iota_a \circ (\mathrm{id}_{V_\mathbf{l}} \otimes \mathbf{r}(T_{\tau^{-1}_{-.d}(i)})) \circ P_a & \text{if } 1 \leq i \leq \ell - 2 \text{ and } p_i = p_{i+1} = 1, \\ \iota_a \circ (\mathrm{id}_{V_\mathbf{l}} \otimes \mathbf{r}(T_{n-1})) \circ P_a & \text{if } i = \ell - 1 \text{ and } p_{\ell-1} = p_\ell = 1,\ a = d^C_+. \\ \iota_a \circ (\mathrm{id}_{V_\mathbf{l}} \otimes \mathbf{r}(T_n)) \circ P_a & \text{if } i = \ell - 1 \text{ and } p_\ell = 1,\ a = d^C_-. \\ \iota_a \circ (\mathrm{id}_{V_\mathbf{l}} \otimes \mathbf{r}(T_n)) \circ P_a & \text{if } i = \ell \text{ and } p_\ell = 1,\ a = d^C_+. \\ \iota_a \circ (\mathrm{id}_{V_\mathbf{l}} \otimes \mathbf{r}(T_{n-1})) \circ P_a & \text{if } i = \ell \text{ and } p_{\ell-1} = p_\ell = 1,\ a = d^C_-, \end{cases} \tag{79}$$

where $\tau_{\pm.d} \in S_{m+n}$ are the ones of (44).

Proof. We can check out this directly in a way similar to that for Proof of Proposition 2.5. □

For $\lambda \in \Lambda_{H_q(W(D_m))}$ and $\mu \in \Lambda_{H_q(W(C_n))}$, we denote $\rho_{H_q(W(D_m)),\lambda} \boxtimes^{CD} \rho_{W(C_n)),\mu}$ by $\rho^{CD}_{q;\lambda,\mu}$ and we denote $C_{V \otimes W}$ for $V = V_{H_q(W(D_m)),\lambda}$ and $W = V_{H_q(W(C_n)),\mu}$ by $C^{CD}_{q;\lambda,\mu}$.

Theorem 2.10. *Let $q \in \mathbb{C}$ and assume that*

$$qP_{W(D_m)}(q)P_{W(C_n)}(q) \neq 0. \tag{80}$$

Then the $\mathbb{C}$-algebra homomorphism

$$\bigoplus_{(\lambda,\mu) \in \Lambda_{H_q(W(D_m))} \times \Lambda_{H_q(W(C_n))}} \rho^{CD}_{q;\lambda,\mu} : \tag{81}$$
$$H_q(\mathfrak{osp}(2m|2n)) \to \bigoplus_{(\lambda,\mu) \in \Lambda_{H_q(W(D_m))} \times \Lambda_{H_q(W(C_n))}} \mathrm{End}_\mathbb{C}(C^{CD}_{q;\lambda,\mu})$$

is an isomorphism. Further we have

$$\dim H_q(\mathfrak{osp}(2m|2n)) = \frac{2^{m+n-1}((m+n-1)!(m+2n))^2}{m!n!}. \tag{82}$$

Moreover $H_q(\mathfrak{osp}(2m|2n))$ is a semisimple $\mathbb{C}$-algebra and a complete set of non-equivalent irreducible representations of $H_q(\mathfrak{osp}(2m|2n))$ is given by $\{\rho^{CD}_{q;\lambda,\mu} | (\lambda, \mu) \in \Lambda_{H_q(W(D_m))} \times \Lambda_{H_q(W(C_n))}\}$.

Proof. Note that $W(D_m) \times W(C_n) = 2^{m+n-1}m!n!$. Then we can prove this theorem in the same way as that for Proof of Theorem 2.8. □

Remark 2.11. Now, by (15), (16) and Theorems 2.6, 2.8 and 2.10, it has turned out that if q is non-zero and not any primitive root of unity, then as a $\mathbb{C}$-algebra, $H_q(\mathfrak{g}) = H_q(W)$ introduced in this section for the Lie superalgebra $\mathfrak{g} = A(m,n)$ or $\mathfrak{osp}(m|2n)$ is very similar to the Iwahori-Hecke algebra $H_q(W_0)$ associated with the Weyl group W_0 of the Lie algebra $\mathfrak{g}(0)$ given as the even part of $\mathfrak{g}$.

Remark 2.12. Assume q to be an element of $\mathbb{C}$ transcendental over $\mathbb{Q}$. Then the $\mathbb{Z}$-subalgebra (with identity) of $\mathbb{C}$ generated by q can also be regarded as the polynomial ring $\mathbb{Z}[q]$ in the variable q over $\mathbb{Z}$. Let W be one of the Coxeter groupoids treated in Subsections 2.5, 2.6 and 2.7. By Lemma 2.2 and (54), (63), (82), one see that $\{f(w)|w \in W \setminus \{0\}\}$ is a $\mathbb{C}$-basis of $H_q(W)$, that is, $H_q(W) = \oplus_{w\in W\setminus\{0\}}\mathbb{C}f(w)$. Define $H_{\mathbb{Z}[q],q}(W)$ to be the $\mathbb{Z}[q]$-submodule generated by $\{f(w)|w \in W \setminus \{0\}\}$. Using Theorem 1.7 and Corollary 1.8, one see that $H_{\mathbb{Z}[q],q}(W)$ is also a $\mathbb{Z}[q]$-subalgebra of $H_q(W)$. Let $\mathbb{A}$ be any commutative ring (with identity). Let ζ be any element of $\mathbb{A}$. Regard $\mathbb{A}$ as a $\mathbb{Z}[q]$-algebra via the $\mathbb{Z}$-algebra homomorphism $\mathbb{Z}[q] \to \mathbb{A}$ that takes q to ζ. Let $H_{\mathbb{A},\zeta}(W)$ be the $\mathbb{A}$-algebra (with identity) defined by $H_{\mathbb{A},\zeta}(W) := H_{\mathbb{Z}[q],q}(W) \otimes_{\mathbb{Z}[q]} \mathbb{A}$. For $X \in H_{\mathbb{Z}[q],q}(W)$, we also denote the element $X \otimes 1$ of $H_{\mathbb{A},\zeta}(W)$ by X. Then $H_{\mathbb{A},\zeta}(W)$ is a free $\mathbb{A}$-module with an $\mathbb{A}$-basis $\{f(w)|w \in W \setminus \{0\}\}$, that is,

$$\mathrm{rank}_{\mathbb{A}} H_{\mathbb{A},\zeta}(W) = |W| - 1. \tag{83}$$

Using Theorem 1.7 and Corollary 1.8 again, one see that $H_{\mathbb{A},\zeta}(W)$ can also be defined by the same generators as (17) and the same relations as (18)-(25) with ζ in place of q.

The same properties as above seem to be true for many Coxeter groupoids, which might be able to be proved in a way similar to that of the proof of [L, Proposition 3.3].

References

[AS1] N. Andruskiewitsch and H.-J. Schneider, *Lifting of quantum linear spaces and pointed Hopf algebras of order p^3*, *J. Algebra* **209** (1998) 658–691.

[AS2] N. Andruskiewitsch and H.-J. Schneider, *On the classification of finite-dimensional pointed Hopf algebras*, Accepted for publication in Ann. Math., [arXiv:math.QA/0502157].

[C] R.W. Carter, Simple groups of Lie type, Pure and Applied Mathematics, Vol. 28. John Wiley & Sons, London-New York-Sydney, 1972.

[CR] C.W. Curtis and I. Reiner, Representation theory of finite groups and associative algebras, Interscience Publishers, a division of John Wiley & Sons, New York-London 1962

[GU] A. Gyoja and K. Uno, *On the semisimplicity of Hecke algabras*, J. Math. Soc. Japan **41** (1989), no. 1, 75–79

[H1] I. Heckenberger, *Classification of arithmetic root systems*, to appear in Adv. Math. **220** (2009) 59-124

[H2] ———, *The Weyl groupoid of a Nichols algebra of diagonal type*, Invent. Math. **164** (2006) 175–188.

[HSTY] I. Heckenberger, F. Spill, A. Torrielli, H. Yamane, *Drinfeld second realization of quantum affine superalgebras of $D^{(1)}(2,1;x)$ via the Weyl groupoid*, *RIMS Kôkyûroku Bessatsu* **B8** (2008) 171-216

[HY] I. Heckenberger and H. Yamane, *A generalization of Coxeter groups, root systems, and Matsumoto's theorem*, Math. Z. **259** (2008) 255-276

[Hu] J. Humphreys, Reflection groups and Coxeter groups. Cambridge Studies in Advanced Mathematics, 29. Cambridge University Press, Cambridge, 1990

[K] V.G. Kac, *Lie superalgebras*, Adv. Math. **26** (1977) 8–96

[L] G. Lusztig, Hecke algebras with unequal parameters. CRM Monograph Series, 18. American Mathematical Society, Providence, RI, 2003.

[S] V. Serganova, *On generalizations of root systems*, Commun. Algebra **24** (1996), 4281–4299.

[SV] A.N. Sergeev and A.P. Veselov, *Grothendieck rings of basic classical Lie superalgebras*, Preprint arXiv:0704.2250

Hiroyuki Yamane, Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Toyonaka 560-0043, Japan

E-mail address: `yamane@ist.osaka-u.ac.jp`